

# QCD factorization beyond leading twist in exclusive processes: $\rho_T$ -meson production

Samuel Wallon

Laboratoire de Physique Théorique  
Université Paris Sud  
Orsay

LPT, january 22th, 2009

in collaboration with

I. V. Anikin (JINR, Dubna), D. Yu. Ivanov (SIM, Novosibirsk), B. Pire (CPhT, Palaiseau) and L. Szymanowski (SINS, Warsaw)

# Outline

- 1 Introduction
  - Exclusive processes at high energy QCD
  - QCD in the perturbative Regge limit
  - Exclusive  $\rho$ -production
- 2 Impact factor for exclusive processes
  - Theoretical motivations
  - $k_T$  factorization
  - Gauge invariance of impact factor
- 3 Collinear factorization
  - Light-Cone Collinear approach
  - Parametrization of vacuum-to-rho-meson matrix elements (DAs)
  - Symmetry properties
  - Equations of motion
  - $n$ -independence
  - A set of independent non-perturbative correlators
- 4 Computation and results
  - Computation of the hard part
  - Results
  - Discussion
- 5 Conclusions

# Introduction

## Exclusive processes at high energy in QCD

- Since a decade, there have been much developpements in hard exclusive processes.
  - form factors, Distribution Amplitudes → Generalized Distribution Amplitudes
  - DVCS → Generalized Parton Distributions, Transition Distribution Amplitudes
- The key tool is the collinear factorization


## Introduction

## Extensions from DIS

- DIS: inclusive process  $\rightarrow$  forward amplitude ( $t = 0$ )

Structure Function


$$= \text{Coefficient Function (hard)} \otimes \text{Parton Distribution Function (soft)}$$



- DVCS: exclusive process  $\rightarrow$  non forward amplitude ( $-t \ll s = W^2$ )

Amplitude

$$= \text{Coefficient Function (hard)} \otimes \text{Generalized Parton Distribution (soft)}$$




## Introduction

## Extensions from GPD


- **Meson production:**  $\gamma$  replaced by  $\rho, \pi, \dots$

$$\text{Amplitude} = \text{GPD (soft)} \otimes \text{CF (hard)} \otimes \text{Distribution Amplitude (soft)}$$



- **Crossed process:**  $s \ll -t$


$$\text{Amplitude} = \text{Coefficient Function (hard)} \otimes \text{Generalized Distribution Amplitude (soft)}$$



## Introduction

## Extensions from GPD

- starting from usual DVCS, one allows **initial hadron  $\neq$  final hadron**  
example:



which can be further extended by replacing the outgoing  $\gamma$  by any hadronic state

$$\text{Amplitude} = \text{Transition Amplitude (soft)} \otimes \text{CF (hard)} \otimes \text{DA (soft)}$$

# Introduction: phenomenology of exclusive processes within collinear factorization

- Experimental tests are possible in **fixed target** experiments

- $e^{\pm}p, \mu^{\pm}p$ : HERA (HERMES), JLab, COMPASS...

as well as in **colliders**, mainly for medium  $s$

- $e^{\pm}p$  colliders: HERA (H1, ZEUS)

- $e^{+}e^{-}$  colliders: LEP, Belle, BaBar, BEPC

- **Collinear factorization** has been proven only for specific cases:  
e.g.:  $\rho_T$  production cannot directly be factorized (appearance of **end point singularities**)  
⇒ improvement needed for a consistent approach of exclusive processes

# QCD in the perturbative Regge limit

- At the same time, **at large  $s$** , the interest for phenomenological tests of **hard Pomeron** and related resummed approaches has become pretty wide:
  - **inclusive** tests (total cross-section) and semi-inclusive tests (diffraction, forward jets, ...)
  - **exclusive** tests (meson production)
- These tests concern all type of collider experiments:
  - $e^\pm p$ : **HERA**: (H1, ZEUS)
  - $p\bar{p}$  and  $pp$ : **TEVATRON** (CDF, D0); **LHC** (CMS, ATLAS, ALICE)
  - $e^+e^-$ : (LEP, ILC)
- These high energy exclusive processes in the perturbative **Regge** limit may provide new ideas when dealing with collinear factorization



# Introduction

## Exclusive $\rho$ -production


Our studies attempt to describe exclusive processes involving the production of  $\rho$ -mesons in diffraction-type experiment. We choose  $t = t_{min}$  for simplicity.

- $\gamma^*(q) + \gamma^*(q') \rightarrow \rho_T(p_1) + \rho(p_2)$  process in  $e^+ e^- \rightarrow e^+ e^- \rho_T(p_1) + \rho(p_2)$  with double tagged lepton at ILC
- $\gamma^*(q) + P \rightarrow \rho_T(p_1) + P$  at HERA

This process was studied by H1 and ZEUS

- the total cross-section strongly **decreases with  $Q^2$**
- dramatic **increase with  $W^2 = s_{\gamma^* P}$**  (transition from soft to hard regime governed by  $Q^2$ )

(from X. Janssen (H1), DIS 2008)



## Introduction

Exclusive  $\rho$ -productionPolarization effects in  $\gamma^* P \rightarrow \rho P$  at HERA

- one can experimentally measure all spin density matrix elements
- at  $t = t_{min}$  one can experimentally distinguish

$$\begin{cases} \gamma_L^* \rightarrow \rho_L : & \text{dominates} & (\text{twist 2 dominance}) \\ \gamma_T^* \rightarrow \rho_T : & \text{sizable} & (\text{twist 3}) \end{cases}$$



- S-channel helicity conservation:

$$\begin{cases} \gamma_L^* \rightarrow \rho_L & (\equiv T_{00}) \\ \gamma_T^* \rightarrow \rho_T, \end{cases}$$

Dominate with respect to all other transitions.

Experimentally,  $\gamma_T^* \rightarrow \rho_T$  is dominated

by  $\gamma_{T(-)}^* \rightarrow \rho_{T(-)}$  and  $\gamma_{T(+)}^* \rightarrow \rho_{T(+)} (\equiv T_{11})$



(from X. Janssen (H1), DIS 2008)

# Introduction

## Exclusive $\rho$ -production

The processes with vector particle such as rho-meson probe deeper into the fine features of QCD.

It deserves theoretical developpement to describe HERA data in its special kinematical range:

- large  $s_{\gamma^* P} \Rightarrow$  small-x effects expected, within  $k_t$ -factorization
- large  $Q^2 \Rightarrow$  hard scale  $\Rightarrow$  perturbative approach and collinear factorization  $\Rightarrow$  the  $\rho$  can be described through its chiral even Distribution Amplitudes

$$\left\{ \begin{array}{ll} \rho_L & \text{twist 2} \\ \rho_T & \text{twist 3} \end{array} \right.$$

The main ingredient is the  $\gamma^* \rightarrow \rho$  impact factor

- For  $\rho_T$ , special care is needed: a pure 2-body description would violate gauge invariance.
- We show that:
  - Including in a consistent way all twist 3 contributions, i.e. 2-body and 3-body correlators, gives a gauge invariant impact factor
  - Our treatment is free of end-point singularities and does not violates the QCD factorization


# Impact factor for exclusive processes

## Theoretical motivations


### QCD in perturbative Regge limit

- In this limit, the dynamics is dominated by gluons (dominance of spin 1 exchange in  $t$  channel)
- BFKL (and extensions: NLL, saturations effects, ...) is expected to dominate with respect to Born order at large relative rapidity.

Born order:




BFKL ladder:



## Impact factor for exclusive processes

 $k_T$  factorization $\gamma^* \gamma^* \rightarrow \rho \rho$  as an example

- Use **Sudakov** decomposition  $k = \alpha p_1 + \beta p_2 + k_\perp$  ( $p_1^2 = p_2^2 = 0$ ,  $2p_1 \cdot p_2 = s$ )
- write  $d^4k = \frac{s}{2} d\alpha d\beta d^2k_\perp$
- $t$ -channel gluons with **non-sense** polarizations ( $\epsilon_{NS}^{up} = \frac{2}{s} p_2$ ,  $\epsilon_{NS}^{down} = \frac{2}{s} p_1$ ) dominate **at large  $s$**



## Impact factor for exclusive processes

 $k_T$  factorization

impact representation



 $\underline{k} = \text{Eucl.} \leftrightarrow k_{\perp} = \text{Mink.}$ 

$$\mathcal{M} = is \int \frac{d^2 \underline{k}}{(2\pi)^2 \underline{k}^2 (\underline{r} - \underline{k})^2} \Phi^{\gamma^*(q_1) \rightarrow \rho(p_1^{\rho})}(\underline{k}, \underline{r} - \underline{k}) \Phi^{\gamma^*(q_2) \rightarrow \rho(p_2^{\rho})}(-\underline{k}, -\underline{r} + \underline{k})$$

The  $\gamma_{L,T}^*(q)g(k_1) \rightarrow \rho_{L,T}g(k_2)$  **impact factor** is normalized as

$$\Phi^{\gamma^* \rightarrow \rho}(\underline{k}^2) = e^{\gamma^* \mu} \frac{1}{2s} \int \frac{d\kappa}{2\pi} \text{Disc}_{\kappa} \mathcal{S}_{\mu}^{\gamma^* g \rightarrow \rho g}(\underline{k}^2),$$

with  $\kappa = (q+k)^2 = \beta s - Q^2 - \underline{k}^2$




# Impact factor for exclusive processes

Gauge invariance within subleading twists

## Gauge invariance

- **QCD gauge invariance** (probes are colorless)  
 $\Rightarrow$  **impact factor should vanish** when  $\underline{k} \rightarrow 0$  or  $\underline{r} - \underline{k} \rightarrow 0$
- In the following we will restrict ourselves to the case  $t = t_{min}$ , i.e. to  $\underline{r} = 0$



$$k_1 = \frac{\kappa + Q^2 + k^2}{s} p_2 + k_\perp$$

$$k_2 = \frac{\kappa + k^2}{s} p_2 + k_\perp,$$

$$k_1^2 = k_2^2 = -\underline{k}^2$$

This kinematics takes into account **skewedness effects** along  $p_2$

$\Rightarrow$  restriction to the transitions  $\left\{ \begin{array}{ll} 0 & \rightarrow 0 \quad (\text{twist } 2) \\ (+ \text{ or } -) & \rightarrow (+ \text{ or } -) \quad (\text{twist } 3) \end{array} \right.$


- At twist 3 level (for  $\gamma_T^* \rightarrow \rho_T$  transition), gauge invariance is a non trivial statement which **requires 2 and 3 body correlators**

# Collinear factorization

## Light-Cone Collinear approach

- The impact factor can be written as

$$\Phi = \int d^4l \dots \text{tr}[\underbrace{H(l \dots)}_{\text{hard part}} \underbrace{S(l \dots)}_{\text{soft part}}]$$



- At the 2-body level:

$$S_{q\bar{q}}(l) = \int d^4z e^{-il \cdot z} \langle \rho(p) | \psi(0) \bar{\psi}(z) | 0 \rangle,$$

- $H$  and  $S$  are related by  $\int d^4l$  and by the **summation over spinor indices**



## Collinear factorization

Light-Cone Collinear approach: 2 steps of factorization (2-body case)

## 1 - Momentum factorization (1)

- Use **Sudakov** decomposition in the form ( $p = p_1$ ,  $n = 2p_2/s \Rightarrow p \cdot n = 1$ )

$$l_\mu = y p_\mu + l_\mu^\perp + (l \cdot p) n_\mu, \quad y = l \cdot n$$

$$\text{scaling:} \quad 1 \quad 1/Q \quad 1/Q^2$$

- decompose  $H(k)$  around the  $p$  direction:

$$H(l) = H(y p) + \left. \frac{\partial H(l)}{\partial l_\alpha} \right|_{l=y p} (l - y p)_\alpha + \dots \quad \text{with } (l - y p)_\alpha \approx l_\alpha^\perp$$

- In **Fourier** space, the **twist 3** term  $l_\alpha^\perp$  turns into a derivative of the **soft term**

$$\Rightarrow \text{one will deal with } \int d^4 z e^{-i l \cdot z} \langle \rho(p) | \psi(0) i \overleftrightarrow{\partial}_{\alpha^\perp} \bar{\psi}(z) | 0 \rangle$$

# Collinear factorization

Light-Cone Collinear approach: **2 steps of factorization** (2-body case)

## 1 - Momentum factorization (2)

- write

$$d^4l \longrightarrow d^4l \delta(\mathbf{y} - l \cdot n) d\mathbf{y}$$

- $\int d^4l \delta(\mathbf{y} - l \cdot n)$  is then absorbed in the soft term:

$$\begin{aligned} (\tilde{S}_{q\bar{q}}, \partial_{\perp} \tilde{S}_{q\bar{q}}) &\equiv \int d^4l \delta(\mathbf{y} - l \cdot n) \int d^4z e^{-il \cdot z} \langle \rho(p) | \psi(0) (1, i \overrightarrow{\partial}_{\perp}) \bar{\psi}(z) | 0 \rangle \\ (\delta(\mathbf{y} - l \cdot n) = \int \frac{d\lambda}{2\pi} e^{-i\lambda(\mathbf{y} - l \cdot n)} \Rightarrow) &= \int \frac{d\lambda}{2\pi} e^{-i\lambda\mathbf{y}} \int d^4z \delta^{(4)}(z - \lambda n) \langle \rho(p) | \psi(0) (1, i \overrightarrow{\partial}_{\perp}) \bar{\psi}(z) | 0 \rangle \\ &= \int \frac{d\lambda}{2\pi} e^{-i\lambda\mathbf{y}} \langle \rho(p) | \psi(0) (1, i \overrightarrow{\partial}_{\perp}) \bar{\psi}(\lambda n) | 0 \rangle \end{aligned}$$


- $\int d\mathbf{y}$  performs the longitudinal momentum factorization

## Collinear factorization

Light-Cone Collinear approach: 2 steps of factorization (2-body case)

## 2 - Spinorial (and color) factorization

- Use Fierz decomposition of the Dirac (and color) matrices  $\psi(0) \bar{\psi}(z)$  and  $\psi(0) i \overleftrightarrow{\partial}_\perp \bar{\psi}(z)$ :



- $\Phi$  has now the simple factorized form:

$$\Phi = \int dx \left\{ \text{tr} [H_{q\bar{q}}(xp) \Gamma] S_{q\bar{q}}^\Gamma(x) + \text{tr} [\partial_\perp H_{q\bar{q}}(xp) \Gamma] \partial_\perp S_{q\bar{q}}^\Gamma(x) \right\}$$

$\Gamma = \gamma^\mu$  and  $\gamma^\mu \gamma^5$  matrices

$$S_{q\bar{q}}^\Gamma(x) = \int \frac{d\lambda}{2\pi} e^{-i\lambda x} \langle \rho(p) | \bar{\psi}(\lambda n) \Gamma \psi(0) | 0 \rangle$$

$$\partial_\perp S_{q\bar{q}}^\Gamma(x) = \int \frac{d\lambda}{2\pi} e^{-i\lambda x} \langle \rho(p) | \bar{\psi}(\lambda n) \Gamma i \overleftrightarrow{\partial}_\perp \psi(0) | 0 \rangle$$

- choose axial gauge condition for gluons, i.e.  $n \cdot A = 0 \Rightarrow$  no Wilson line

# Collinear factorization

Light-Cone Collinear approach: 2 steps of factorization (3-body case)

## Factorization of 3-body contributions

- 3-body contributions start at **genuine twist 3**  
 ⇒ no need for **Taylor** expansion
- Momentum factorization goes in the same way as for 2-body case
- Spinorial (and color) factorization is similar:



# Collinear factorization

Parametrization of vacuum-to-rho-meson matrix elements (DAs): 2-body correlators

2-body **non-local** correlators

$\rho_L$

**twist 2**

- vector correlator

$$\langle \rho(p) | \bar{\psi}(z) \gamma_\mu \psi(0) | 0 \rangle \stackrel{\mathcal{F}}{=} m_\rho f_\rho \left[ \varphi_1(y) (e^* \cdot n) p_\mu + \varphi_3(y) e_\mu^{*T} \right]$$

$\rho_T$

**kinematical twist 3 (WW)**

**genuine twist 3**

**genuine + kinematical twist 3**

- axial correlator

$$\langle \rho(p) | \bar{\psi}(z) \gamma_5 \gamma_\mu \psi(0) | 0 \rangle \stackrel{\mathcal{F}}{=} m_\rho f_\rho i \varphi_A(y) \varepsilon_{\mu\lambda\beta\delta} e_\lambda^{*T} p_\beta n_\delta$$

- vector correlator with transverse derivative

$$\langle \rho(p) | \bar{\psi}(z) \gamma_\mu i \overleftrightarrow{\partial}_\alpha^\perp \psi(0) | 0 \rangle \stackrel{\mathcal{F}}{=} m_\rho f_\rho \varphi_1^T(y) p_\mu e_\alpha^{*T}$$

- axial correlator with transverse derivative

$$\langle \rho(p) | \bar{\psi}(z) \gamma_5 \gamma_\mu i \overleftrightarrow{\partial}_\alpha^\perp \psi(0) | 0 \rangle \stackrel{\mathcal{F}}{=} m_\rho f_\rho i \varphi_A^T(y) p_\mu \varepsilon_{\alpha\lambda\beta\delta} e_\lambda^{*T} p_\beta n_\delta,$$

where  $y$  ( $\bar{y} \equiv 1 - y$ ) = momentum fraction along  $p \equiv p_1$  of the quark (antiquark) and

$$\stackrel{\mathcal{F}}{=} \int_0^1 dy \exp[i y p \cdot z], \text{ with } z = \lambda n$$

⇒ 5 2-body DAs

# Collinear factorization

Parametrization of vacuum-to-rho-meson matrix elements: 3-body correlators

## 3-body non-local correlators

genuine twist 3

- vector correlator

$$\langle \rho(p) | \bar{\psi}(z_1) \gamma_\mu g A_\alpha^T(z_2) \psi(0) | 0 \rangle \stackrel{\mathcal{F}_2}{\equiv} m_\rho f_3^V B(y_1, y_2) p_\mu e_\alpha^{*T},$$

- axial correlator

$$\langle \rho(p) | \bar{\psi}(z_1) \gamma_5 \gamma_\mu g A_\alpha^T(z_2) \psi(0) | 0 \rangle \stackrel{\mathcal{F}_2}{\equiv} m_\rho f_3^A i D(y_1, y_2) p_\mu \varepsilon_{\alpha\lambda\beta\delta} e_\lambda^{*T} p_\beta n_\delta,$$

where  $y_1, \bar{y}_2, y_2 - y_1 =$  quark, antiquark, gluon momentum fraction

and  $\stackrel{\mathcal{F}_2}{\equiv} \int_0^1 dy_1 \int_0^1 dy_2 \exp[i y_1 p \cdot z_1 + i(y_2 - y_1) p \cdot z_2]$ , with  $z_{1,2} = \lambda n$

$\Rightarrow$  2 3-body DAs

# Collinear factorization

## Symmetry properties

From **C-conjugation** on the previous correlators, one gets:

- 2-body correlators:

$$\varphi_1(y) = \varphi_1(1-y)$$

$$\varphi_3(y) = \varphi_3(1-y)$$

$$\varphi_A(y) = -\varphi_A(1-y)$$

$$\varphi_1^T(y) = -\varphi_1^T(1-y)$$

$$\varphi_A^T(y) = \varphi_A^T(1-y)$$

- 3-body correlators:

$$B(y_1, y_2) = -B(1-y_2, 1-y_1)$$

$$D(y_1, y_2) = D(1-y_2, 1-y_1)$$

# Collinear factorization

## Equations of motion

### Equations of motion

twist 2  
kinematical twist 3 (WW)  
genuine twist 3  
genuine + kinematical twist 3

- Dirac equation leads to

$$\langle i(\overrightarrow{\mathcal{D}}(0)\psi(0))_\alpha \bar{\psi}_\beta(z) \rangle = 0 \quad (i \overrightarrow{D}_\mu = i \overrightarrow{\partial}_\mu + A_\mu)$$

- Apply the Fierz decomposition to the above 2 and 3-body correlators

$$-\langle \psi(x) \bar{\psi}(z) \rangle = \frac{1}{4} \langle \bar{\psi}(z) \gamma_\mu \psi(x) \rangle \gamma_\mu + \frac{1}{4} \langle \bar{\psi}(z) \gamma_5 \gamma_\mu \psi(x) \rangle \gamma_\mu \gamma_5.$$

- $\Rightarrow$  Equation of motion:

$$\begin{aligned} \bar{y}_1 \varphi_3(y_1) + \bar{y}_1 \varphi_A(y_1) + \varphi_1^T(y_1) + \varphi_A^T(y_1) \\ + \int dy_2 \left[ \zeta_3^V B(y_1, y_2) + \zeta_3^A D(y_1, y_2) \right] = 0 \end{aligned} \quad (1)$$

- In WW approximation: genuine twist 3 = 0 i.e. B=D=0

$$\begin{cases} \varphi_A^T(y) = \frac{1}{2}[(y - \bar{y}) \varphi_3^{WW}(y) - \varphi_3^{WW}(y)] \\ \varphi_1^T(y) = \frac{1}{2}[(y - \bar{y}) \varphi_3^{WW}(y) - \varphi_A^{WW}(y)] \end{cases}$$



## Collinear factorization

 $n$ -independence

## A minimal set of DAs

- The non-perturbative correlators cannot be obtained from perturbative QCD (!)
- one should reduce them to a minimal set before using any model
- this can be achieved by using an additional condition:  
independency of the full amplitude with respect to the gauge fixing vector  $n$

⇒ we prove that 3 independent Distribution Amplitudes are needed:

$\phi_1(y)$  ← 2 body twist 2 correlator

$B(y_1, y_2)$  ← 3 body genuine twist 3 vector correlator

$D(y_1, y_2)$  ← 3 body genuine twist 3 axial correlator

## Collinear factorization


 $n$ -independence $n$ -independence in practice

- $\rho_T$  polarization:  $e_\mu^{*T} = e_\mu^* - p_\mu e^* \cdot n$  keeping  $n \cdot p = 1$
- for the full factorized amplitude:

$$A = H \otimes S \quad \frac{dA}{dn^\mu} = 0, \quad \text{where} \quad \frac{d}{dn^\mu} = \frac{\partial}{\partial n^\mu} + e_\mu^* \frac{\partial}{\partial (e^* \cdot n)}$$

- rewrite hard terms in one single form, of 2-body type: use Ward identities  
Example: hard 3-body  $\rightarrow$  hard 2-body

$$\text{tr} [H_{3\rho}(y_1, y_2) p^\rho \not{p}] B(y_1, y_2) = \frac{1}{y_1 - y_2} (\text{tr} [H_2(y_1) \not{p}] - \text{tr} [H_2(y_2) \not{p}]) B(y_1, y_2),$$



- thus, symbolically,

$$\frac{dS}{dn^\mu} = 0$$

## Collinear factorization

 $n$ -independenceConstraints from  $n$ -independence

twist 2

kinematical twist 3 (WW)

genuine twist 3

genuine + kinematical twist 3

- vector correlators

$$\frac{d}{dy_1} \varphi_1^T(y_1) = -\varphi_1(y_1) + \varphi_3(y_1)$$

$$-\zeta_3^V \int_0^1 \frac{dy_2}{y_2 - y_1} (B(y_1, y_2) + B(y_2, y_1))$$

- axial correlators

$$\frac{d}{dy_1} \varphi_A^T(y_1) = \varphi_A(y_1) - \zeta_3^A \int_0^1 \frac{dy_2}{y_2 - y_1} (D(y_1, y_2) + D(y_2, y_1))$$

# Collinear factorization

A set of independent non-perturbative correlators

## Solution

twist 2  
 kinematical twist 3 (WW)  
 genuine twist 3  
 genuine + kinematical twist 3

- the set of 4 equations (2 EOM + 2  $n$ -independence relations) can be solved analytically
- 7  $\rightarrow$  3 independent DAs

# Collinear factorization

A set of independent non-perturbative correlators

Solution in WW approximation

twist 2

kinematical twist 3 (WW)

genuine twist 3

genuine + kinematical twist 3

WW:  $B = D = 0$

$$\varphi_{3/A}^{WW}(y) = \frac{1}{2} \left[ \int_0^y \frac{dv}{\bar{v}} \varphi_1(v) \pm \int_y^1 \frac{dv}{v} \varphi_1(v) \right]$$

$$\varphi_{1/A}^{TWW}(y) = \frac{1}{2} \left[ -\bar{y} \int_0^y \frac{dv}{\bar{v}} \varphi_1(v) \pm y \int_y^1 \frac{dv}{v} \varphi_1(v) \right]$$

## Collinear factorization

A set of independent non-perturbative correlators

Solutions: genuine twist-3 vector part

twist 2

kinematical twist 3 (WW)

genuine twist 3

genuine + kinematical twist 3

- vector 2-body DAs for  $\rho_T$ :  $\varphi_3(y)$  and  $\varphi_1^T(y)$

- $\varphi_3(y) = \varphi_3^{WW}(y) + \varphi_3^{gen}(y)$

$$\varphi_3^{gen}(y) = -\frac{1}{2} \int_y^1 \frac{du}{u} \left[ \int_0^u dy_2 \frac{d}{du} (\zeta_3^V B - \zeta_3^A D)(y_2, u) - \int_u^1 \frac{dy_2}{y_2 - u} (\zeta_3^V B - \zeta_3^A D)(u, y_2) \right. \\ \left. - \int_0^u \frac{dy_2}{y_2 - u} (\zeta_3^V B - \zeta_3^A D)(y_2, u) \right] \\ - \frac{1}{2} \int_0^{y_1} \frac{du}{\bar{u}} \left[ \int_u^1 dy_2 \frac{d}{du} (\zeta_3^V B + \zeta_3^A D)(u, y_2) - \int_u^1 \frac{dy_2}{y_2 - u} (\zeta_3^V B + \zeta_3^A D)(u, y_2) \right. \\ \left. - \int_0^u \frac{dy_2}{y_2 - u} (\zeta_3^V B + \zeta_3^A D)(y_2, u) \right]$$

- $\varphi_1^T(y) = \varphi_1^{TWW}(y) + \varphi_1^{gen}(y)$

$$\varphi_1^{Tgen}(y) = \int_0^y du \varphi_3^{gen}(u) - \zeta_3^V \int_0^y dy_1 \int_y^1 dy_2 \frac{B(y_1, y_2)}{y_2 - y_1}$$

# Collinear factorization

A set of independent non-perturbative correlators

Solutions: genuine twist-3 axial part

twist 2

kinematical twist 3 (WW)

genuine twist 3

genuine + kinematical twist 3

- axial 2-body DAs for  $\rho_T$ :  $\varphi_A(y)$  and  $\varphi_A^T(y)$
- $\varphi_A(y) = \varphi_A^{WW}(y) + \varphi_A^{gen}(y)$
- $\varphi_A^T(y) = \varphi_A^{TWW}(y) + \varphi_A^{gen}(y)$
- the corresponding expressions for  $\varphi_A^{gen}(y)$  and  $\varphi_A^{Tgen}(y)$  are obtained by the substitutions:



$$\begin{array}{ccc}
 \varphi_A^{gen}(y) & \longleftrightarrow & \varphi_3^{gen}(y) \\
 \zeta_3^V B \leftrightarrow \zeta_3^A D & & \\
 \\
 \varphi_A^{Tgen}(y) & \longleftrightarrow & \varphi_1^{Tgen}(y) \\
 \zeta_3^V B \leftrightarrow \zeta_3^A D & & 
 \end{array}$$

# Computation and results

## Computation of the hard part

### 2-body diagrams

- without derivative







twist 2  $(\gamma_L^* \rightarrow \rho_L)$

twist 3  $(\gamma_T^* \rightarrow \rho_T)$

- practical trick for computing  $\partial_\perp H$  : use the **Ward identity**

$$\frac{\partial}{\partial p_\mu} \rightarrow p = \rightarrow p \bullet \gamma^\mu \rightarrow p$$

$$\text{where } \rightarrow p = \frac{1}{m - \not{p} - i\epsilon}$$






# Computation and results


## Computation of the hard part

### 3-body diagrams

- “abelian” type



- “non-abelian” type



# Computation and results

Recall:  $\gamma_L^* \rightarrow \rho_L$  impact factor

$\gamma_L^* \rightarrow \rho_L$  impact factor

$$\Phi^{\gamma_L^* \rightarrow \rho_L}(\underline{k}^2) = \frac{2 e g^2 f_\rho}{Q} \frac{\delta^{ab}}{2 N_c} \int dy \varphi_1(y) \frac{\underline{k}^2}{y \bar{y} Q^2 + \underline{k}^2}$$

pure twist 2 scaling

# Computation and results

Results:  $\gamma_T^* \rightarrow \rho_T$  impact factor

$\gamma_T^* \rightarrow \rho_T$  impact factor:

Spin Non-Flip/Flip separation appears

$$\Phi^{\gamma_T^* \rightarrow \rho_T}(\underline{k}^2) = \Phi_{n.f.}^{\gamma_T^* \rightarrow \rho_T}(\underline{k}^2) T_{n.f.} + \Phi_{f.}^{\gamma_T^* \rightarrow \rho_T}(\underline{k}^2) T_f.$$

where

$$T_{n.f.} = -(e_\gamma \cdot e^*) \quad \text{and} \quad T_f = \frac{(e_\gamma \cdot k)(e^* k)}{\underline{k}^2} + \frac{(e_\gamma \cdot e^*)}{2}$$

non-flip transitions  $\left\{ \begin{array}{l} + \rightarrow + \\ - \rightarrow - \end{array} \right.$

flip transitions  $\left\{ \begin{array}{l} + \rightarrow - \\ - \rightarrow + \end{array} \right.$

# Computation and results

Results:  $\gamma_T^* \rightarrow \rho_T$  impact factor

pure twist 3 scaling

$$\begin{aligned} & \Phi_{n.f.}^{\gamma_T^* \rightarrow \rho_T}(\underline{k}^2) \\ &= -\frac{e g^2 m_\rho f_\rho}{2\sqrt{2} Q^2} \frac{\delta^{ab}}{2 N_c} \left\{ -2 \int dy_1 \frac{(\underline{k}^2 + 2 Q^2 y_1 (1 - y_1)) \underline{k}^2}{y_1 (1 - y_1) (\underline{k}^2 + Q^2 y_1 (1 - y_1))^2} \left[ (2y_1 - 1) \varphi_1^T(y_1) + \varphi_A^T(y_1) \right] \right. \\ &+ 2 \int dy_1 dy_2 \left[ \zeta_3^V B(y_1, y_2) - \zeta_3^A D(y_1, y_2) \right] \frac{y_1 (1 - y_1) \underline{k}^2}{\underline{k}^2 + Q^2 y_1 (1 - y_1)} \left[ \frac{(2 - N_c/C_F) Q^2}{\underline{k}^2 (y_1 - y_2 + 1) + Q^2 y_1 (1 - y_2)} \right. \\ &- \left. \frac{N_c}{C_F} \frac{Q^2}{y_2 \underline{k}^2 + Q^2 y_1 (y_2 - y_1)} \right] - 2 \int dy_1 dy_2 \left[ \zeta_3^V B(y_1, y_2) + \zeta_3^A D(y_1, y_2) \right] \left[ \frac{2 + N_c/C_F}{1 - y_1} \right. \\ &+ \frac{y_1 Q^2}{\underline{k}^2 + Q^2 y_1 (1 - y_1)} \left( \frac{(2 - N_c/C_F) y_1 \underline{k}^2}{\underline{k}^2 (y_1 - y_2 + 1) + Q^2 y_1 (1 - y_2)} - 2 \right) \\ &\left. \left. + \frac{N_c (y_1 - y_2) (1 - y_2)}{C_F (1 - y_1)} \frac{Q^2}{\underline{k}^2 (1 - y_1) + Q^2 (y_2 - y_1) (1 - y_2)} \right] \right\} \end{aligned}$$

and

$$\begin{aligned} & \Phi_{f.}^{\gamma_T^* \rightarrow \rho_T}(\underline{k}^2) = -\frac{e g^2 m_\rho f_\rho}{2\sqrt{2} Q^2} \frac{\delta^{ab}}{2 N_c} \left\{ 4 \int dy_1 \frac{\underline{k}^2 Q^2}{(\underline{k}^2 + Q^2 y_1 (1 - y_1))^2} \left[ \varphi_A^T(y_1) - (2y_1 - 1) \varphi_1^T(y_1) \right] \right. \\ &- 4 \int dy_1 dy_2 \frac{y_1 \underline{k}^2}{\underline{k}^2 + Q^2 y_1 (1 - y_1)} \left[ \zeta_3^A D(y_1, y_2) (-y_1 + y_2 - 1) + \zeta_3^V B(y_1, y_2) (y_1 + y_2 - 1) \right] \\ &\left. \times \left[ \frac{(2 - N_c/C_F) Q^2}{\underline{k}^2 (y_1 - y_2 + 1) + Q^2 y_1 (1 - y_2)} - \frac{N_c}{C_F} \frac{Q^2}{y_2 \underline{k}^2 + Q^2 y_1 (y_2 - y_1)} \right] \right\} \end{aligned}$$

# Computation and results

Results:  $\gamma_T^* \rightarrow \rho_T$  impact factor

## WW limit

- WW limit: keep only **twist 2** + **kinematical twist 3** terms (i.e  $B = D = 0$ )
- The only remaining contributions come from the two-body correlators
- non-flip transition

$$\Phi_{n.f.}^{\gamma_T^* \rightarrow \rho_T}(\underline{k}^2) = \frac{-e m_\rho f_\rho}{2\sqrt{2} Q^2} \frac{\delta^{ab}}{2 N_c} \int_0^1 dy \left\{ \frac{(y - \bar{y}) \varphi_1^{TWW}(y) + 2y\bar{y} \varphi_3^{WW}(y) + \varphi_A^{TWW}(y)}{y\bar{y}} - \frac{2\underline{k}^2 (\underline{k}^2 + 2Q^2 y\bar{y}) \left( (y - \bar{y}) \varphi_1^{TWW}(y) + \varphi_A^{TWW}(y) \right)}{y\bar{y} (\underline{k}^2 + Q^2 y(1-y))^2} \right\}$$

which simplifies, using equation of motion:

$$\int dy \left[ (y - \bar{y}) \varphi_1^{TWW}(y) + 2y\bar{y} \varphi_3^{WW}(y) + \varphi_A^{TWW}(y) \right] = 0$$

$$\Phi_{n.f.}^{\gamma_T^* \rightarrow \rho_T}(\underline{k}^2) = \frac{e m_\rho f_\rho}{\sqrt{2} Q^2} \frac{\delta^{ab}}{2 N_c} \int_0^1 dy \frac{2\underline{k}^2 (\underline{k}^2 + 2Q^2 y\bar{y})}{y\bar{y} (\underline{k}^2 + Q^2 y\bar{y})^2} \left[ (2y - 1) \varphi_1^{TWW}(y) + \varphi_A^{TWW}(y) \right].$$

- flip transition:

$$\Phi_{n.f.}^{\gamma_T^* \rightarrow \rho_T}(\underline{k}^2) = -\frac{e m_\rho f_\rho}{\sqrt{2} Q^2} \frac{\delta^{ab}}{2 N_c} \int_0^1 \frac{2\underline{k}^2 Q^2}{(\underline{k}^2 + Q^2 y\bar{y})^2} \left[ (1 - 2y) \varphi_1^{TWW}(y) + \varphi_A^{TWW}(y) \right].$$

# Computation and results

Discussion: [gauge invariance](#)

- The obtained results are gauge invariant:

$$\Phi^{\gamma_T^* \rightarrow \rho_T} \rightarrow 0 \quad \text{when} \quad \underline{k} \rightarrow 0$$

- this is straightforward in the WW limit
- at the full twist 3 order:
  - the  $C_F$  part of the abelian 3-body contribution cancels the 2-body contribution **after using the equation of motion**
  - the  $N_c$  part of the abelian 3-body contribution cancels the 3-body non-abelian contribution
  - thus  $\gamma_T^* \rightarrow \rho_T$  impact factor is **gauge-invariant only provided the 3-body contributions have been taken into account**

# Computation and results

Discussion: [consistence with factorization](#)

- Our results are free of end-point singularities, in both WW approximation and full twist-3 order calculation:
  - the flip contribution obviously does not have any end-point singularity because of the  $\underline{k}^2$  which regulates them
  - the potential end-point singularity for the non-flip contribution is spurious since  $\varphi_A^T(y)$ ,  $\varphi_1^T(y)$  vanishes at  $y = 0, 1$  as well as  $B(y_1, y_2)$  and  $D(y_1, y_2)$ .

# Conclusions

1

- We have performed a full up to twist 3 computation of the  $\gamma^* \rightarrow \rho$  impact factor, in the  $t = t_{min}$  limit.
- Our result respects gauge invariance. This is achieved only after including 2 and 3 body correlators.
- It is free of end-point singularities (this should be contrasted with standard collinear treatment, at moderate  $s$ , where  $k_T$ -factorization is NOT applicable: see Mankiewicz-Piller).
- Phenomenological applications will be done in the near future.
- In this talk we relied on the Light-Cone Collinear approach (Ellis + Furmanski + Petronzio; Efremov + Teryaev; Anikin + Teryaev), which is non-covariant, but very efficient for practical computations.
- This Light-Cone Collinear approach is systematic, and can be extended to any process, including higher twist effects (but does not preclude potential end-point singularities)



# Conclusions

2

- Comparison with a fully **covariant approach** by **Ball+Braun et al**:  
The dictionary between the two approaches within a full twist 3 treatment is now established:

$$B(y_1, y_2) = -\frac{V(y_1, 1 - y_2, y_2 - y_1)}{y_2 - y_1},$$

$$D(y_1, y_2) = \frac{A(y_1, 1 - y_2, y_2 - y_1)}{y_2 - y_1}$$

$$\varphi_1(y) = f_\rho m_\rho \phi_{\parallel}(y)$$

$$\varphi_3(y) = f_\rho m_\rho g^{(v)}(y),$$

$$\varphi_A(y) = -\frac{1}{4} f_\rho m_\rho \frac{\partial g^{(a)}(y)}{\partial y}$$

- We also performed calculations of the same impact factor within the **covariant approach** by **Ball+Braun et al**: calculations proceed in quite different way : eg. no  $\varphi_{1,A}^T$ -DAs but **Wilson** line effects are important !!  
We got a full agreement with our approach